


In the claims:

1. (previously presented) A process for the preparation of acylphosphanes of formula I



(I), wherein

n and **m** are each independently of the other 1 or 2;



R₁, if **n = 1**, is

C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms, phenyl-C₁-C₄alkyl, C₂-C₈alkenyl, phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or a 5- or 6-membered O-, S- or N-containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkylthio, C₁-C₈alkoxy and/or -N(R₈)₂;


R₁, if **n = 2**, is

C₁-C₁₈alkylene, C₂-C₁₈alkylene which is interrupted by one or several non-successive O atoms; or R₁ is C₁-C₆alkylene which is substituted by C₁-C₄alkoxy, phenyl, C₁-C₄alkylphenyl, phenyl-C₁-C₄alkyl or C₁-C₆alkoxyphenyl; or R₁ is phenylene or xylylene, which radicals are unsubstituted or substituted by

one to three C₁-C₄alkyl and/or C₁-C₄alkoxy, or R₁ is a -CH₂CH=CHCH₂-, -CH₂-C≡C-CH₂-



or



group;

R₂ is C₁-C₁₈alkyl, C₃-C₁₂cycloalkyl, C₂-C₁₈alkenyl, phenyl-C₁-C₄alkyl, phenyl, naphthyl, biphenyl or a 5- or 6-membered O-, S- or N-containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkoxy and/or C₁-C₈alkylthio;

R₃ is C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms or which is interrupted by -CO-, -COO-, -OCO-, -OCOO-, -CO-N(R₉)-, -N(R₉)-CO-, -N(R₉)-CO-N(R₉)-, -N(R₉)-COO-; C₁-C₁₈alkyl substituted by -OR₁₀, -OCO-R₁₀, -COO-R₁₀, -N(R₉)-CO-R₁₀, -CO-N(R₉)-R₁₀, -C(R₁₁)=C(R₁₂)-CO-OR₁₀ or -C(R₁₁)=C(R₁₂)-phenyl;

C₂-C₁₂alkenyl or C₂-C₁₂alkenyl which is interrupted by one or several non-successive O atoms; phenyl-C₁-C₄alkyl, phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or a 5- or 6-membered O-, S- or N-containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkylthio C₁-C₈alkoxy and/or -N(R₈)₂; or R₃ is -CO-OR₉ or -CO-N(R₉)₂;

Q is a single bond, CR₆R₇, -O- or -S-;

R₄ and R₅ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy;

R₆ and R₇ are each independently of the other hydrogen or C₁-C₄alkyl;

R₈ is C₁-C₁₈ alkyl, C₂-C₁₈ alkyl which is interrupted by one or several non-successive O-atoms; or -N(R₈)₂ forms a 5- or 6-membered O-, S- or N-containing heterocyclic ring;

R₉ is hydrogen, C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms, C₃-C₁₂-cycloalkyl, C₂-C₁₈-alkenyl, phenyl-C₁-C₄-alkyl, phenyl, naphthyl, pyridyl, the radicals phenyl, naphthyl or pyridyl being unsubstituted or substituted by one to five C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio and/or halogen; or -N(R₉)₂ forms a 5- or 6-membered O-, S- or N-containing heterocyclic ring;


R₁₀ is C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O-atoms, C₃-C₁₂-cycloalkyl, phenyl-C₁-C₄-alkyl, C₂-C₁₈-alkenyl, phenyl, naphthyl or biphenyl, the radicals phenyl-C₁-C₄-alkyl, phenyl, naphthyl or biphenyl being unsubstituted or substituted by one to five C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio and/or halogen;

R₁₁ is hydrogen or C₁-C₄-alkyl;

R₁₂ is hydrogen or C₁-C₄-alkyl;

by


(1) reacting a phosphorous halide of formula IIa or a phosphorous halide oxide of formula IIb or a phosphorous halide sulfide of formula IIc



wherein R₁, R₃, n and m have the meaning cited above and Hal is F, Cl, Br or I;

with an alkali metal in a solvent (**metallation**) in the presence of a proton source (**reduction**);

(2) subsequent reaction with m acid halides of formula III



wherein R₂, Hal and m have the meaning cited above.

2. **(currently amended)** A process according to claim 1, wherein in step (1) the metallation is carried out by reacting a compound of the formula IIa, IIb, or IIc with ~~an alkali metal in a solvent, whereby a metallized phosphanide of the formula V~~



~~is formed together with cyclic phosphanes (R₄P)_n, n ≥ 3 as intermediates, wherein~~

~~Me is lithium, sodium or potassium or magnesium in combination with lithium in a solvent, and~~

~~R₄ is as defined in claim 1; and~~

~~wherein the reduction is carried out by reacting the intermediate V and/or (R₄P)_n, n ≥ 3 with a proton source.~~

3. **(previously presented)** A process according to claim 2, wherein

the alkali metal is sodium;

the proton source is selected from the group consisting of sterically hindered alcohols, trialkylamine hydrohalogenes, bisarylamines, malono nitrile, malonic acid esters, amidine hydrohalogene and carboxylic acids;

the solvent is one or more compounds selected from the group consisting of benzene, toluene, o-, m- or p-xylene, mesitylene, ethylbenzene, diphenylethane, 1,2,3,4-tetrahydronaphthaline (tetraline) and isopropylbenzene (cumol); and

the reaction temperature of step (1) is in the range from -20°C to +160°C.


4. **(previously presented)** A process according to claim 3, wherein the sterically hindered alcohol is selected from the group consisting of secondary and tertiary C₃-C₁₈alcohols.

5. **(previously presented)** A process according to claim 1, wherein metallation is carried out in the presence of catalytic amounts of alkali or earth alkali hydroxides or of Na, K or Li alcoholates or of alcohols.

6. **(previously presented)** A process according to claim 1, wherein the metallation and reduction step is carried out in the presence of an activator.

7. **(previously presented)** A process according to claim 6, wherein the activator is an amine selected from the group consisting of triethylamine, tributylamine, piperidine, morpholine, N-methylpiperidine, N-methyl morpholine and polyamines.

8. **(previously presented)** A process according to claim 1 for the preparation of **monoacylphosphanes** of the formula I'




(1) reacting organic phosphorus halides of formula II'



with an alkali metal in a solvent in the presence of a proton source; and either


(2a) subsequent reaction with an acid halide of formula III'



followed by the reaction with an electrophilic compound $\text{R}_3\text{-Hal}$, or


(2b) subsequent reaction with with an electrophilic compound $\text{R}_3\text{-Hal}$

followed by the reaction with an acid halide of formula III'



wherein R_1 , R_2 and R_3 and Hal are as defined in claim 1.

9. **(previously presented)** A process according to claim 1 for the preparation of **symmetric bisacylphosphanes** of the formula I''




(1) reacting organic phosphorus halides of formula II''




with an alkali metal in a solvent in the presence of a proton source;

(2) subsequent reaction with an acid halide of formula III''



wherein R_1 and R_2 and Hal are as defined in claim 1.

10. **(original)** A process according to claim 1 for the preparation of **unsymmetric bisacylphosphanes** of the formula I'''




(1) reacting organic phosphorus halides of formula II''




with an alkali metal in a solvent in the presence of a proton source;

(2) subsequent reaction with an acid halide of formula III''



(3) subsequent reaction with a second acid halide III'''



wherein


R_1 is as defined in claim 1 and

R_2 and R_2' independently of one another are as defined in claim 1 under R_2 with the proviso that R_2 is not equal R_2' ,

Hal is as defined in claim 1.

11. (cancelled)

12. (previously presented) A process according to claim 1 for the preparation of mono acylated phosphanes of the formula VI and VI'





(1) reacting organic phosphorus halides of formula II"




with an alkali metal in a solvent in the presence of a proton source;

(2) subsequent reaction with an acid halide of formula III" or III'"



wherein R_1 , R_2 are as defined in claim 1, R_2' is as defined in claim 1 under R_2 with the proviso that R_2 is not equal R_2' and Me is Li, Na, K or Mg in combination with Li.

13. (currently amended) A process according to claim 1, further comprising an additional step of oxidizing the acylphosphane of formula (I) to prepare acylphosphane oxides or reacting the acylphosphane of formula (I) with sulfur to prepare ~~acylphosphane oxides and~~ acylphosphane sulfides of formula IV



(IV), wherein

R_1 , R_2 , R_3 , n and m are as defined in claim 1, and Z is O or S.

14. **(previously presented)** A process according to claim 4, wherein the secondary and tertiary C₃-C₁₈alcohols are selected from the group consisting of t-butanol, t-amyl-alcohol, 3-methyl-3-pentanol, 3-ethyl-3-pentanol, triphenylmethanol, 3,7-dimethyl-3-octanol, 2-methyl-1-phenyl-2-propanol, 2-methyl-4-phenyl-2-butanol, fenchyl alcohol, 2,4-dimethyl-3-pentanol, 1-dimethylamino-2-propanol or hexylene glycol.

15. **(previously presented)** A process according to claim 5, wherein metallation is carried out in the presence of catalytic amounts of Na, K or Li sterically hindered alcoholates or sterically hindered alcohols.

16. **(previously presented)** A process according to claim 7, wherein the polyamine is N,N,N',N'-tetramethylethylenediamine.

17. **(previously presented)** A process according to claim 13, wherein a monoacylphosphane of

formula I'
$$\begin{array}{c} \text{R}_3 \\ | \\ \text{R}_1-\text{P}-\text{C}-\text{R}_2 \\ || \\ \text{O} \end{array}$$
 (I'), is oxidized or reacted sulfur.


18. **(previously presented)** A process according to claim 13, wherein a symmetric bisacylphosphane

of formula I''
$$\begin{array}{c} \text{O} \quad \quad \text{O} \\ || \quad \quad || \\ \text{R}_2-\text{C}-\text{P}-\text{C}-\text{R}_2 \\ | \\ \text{R}_1 \end{array}$$
 (I''), is oxidized or reacted sulfur.

19. **(previously presented)** A process according to claim 13, wherein an unsymmetric

bisacylphosphane of formula I'''
$$\begin{array}{c} \text{O} \quad \quad \text{O} \\ || \quad \quad || \\ \text{R}_2'-\text{C}-\text{P}-\text{C}-\text{R}_2 \\ | \\ \text{R}_1 \end{array}$$
 (I'''), is oxidized or reacted sulfur.

20. (new) A process for the preparation of acylphosphanes of formula I



(I), wherein





n and **m** are each independently of the other 1 or 2;

R₁, if **n = 1**, is

C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms, phenyl-C₁-C₄alkyl, C₂-C₈alkenyl, phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or a 5- or 6-membered O-, S- or N-containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkylthio, C₁-C₈alkoxy and/or -N(R₈)₂;

R₁, if **n = 2**, is

C₁-C₁₈alkylene, C₂-C₁₈alkylene which is interrupted by one or several non-successive O atoms; or R₁ is C₁-C₆alkylene which is substituted by C₁-C₄alkoxy, phenyl, C₁-C₄alkylphenyl, phenyl-C₁-C₄alkyl or C₁-C₆alkoxyphenyl; or R₁ is phenylene or xylylene, which radicals are unsubstituted or substituted by one to three C₁-C₄alkyl and/or C₁-C₄alkoxy, or R₁ is a -CH₂CH=CHCH₂-,



or

group;

R₂ is C₁-C₁₈alkyl, C₃-C₁₂cycloalkyl, C₂-C₁₈alkenyl, phenyl-C₁-C₄alkyl, phenyl, naphthyl, biphenyl or a 5- or 6-membered O-, S- or N-containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkoxy and/or C₁-C₈alkylthio;

R₃ is C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms or which is interrupted by -CO-, -COO-, -OCO-, -OCOO-, -CO-N(R₉)-, -N(R₉)-CO-, -N(R₉)-CO-N(R₉)-, -N(R₉)-COO-; C₁-C₁₈alkyl substituted by -OR₁₀, -OCO-R₁₀, -COO-R₁₀, -N(R₉)-CO-R₁₀, -CO-N(R₉)-R₁₀, -C(R₁₁)=C(R₁₂)-CO-OR₁₀ or -C(R₁₁)=C(R₁₂)-phenyl;

C₂-C₁₂alkenyl or C₂-C₁₂alkenyl which is interrupted by one or several non-successive O atoms; phenyl-C₁-C₄alkyl, phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or a 5- or 6-membered O-, S- or N-

containing heterocyclic ring, the radicals phenyl, naphthyl, biphenyl, C₅-C₁₂cycloalkyl or the 5- or 6-membered O-, S- or N-containing heterocyclic ring being unsubstituted or substituted by one to five halogen, C₁-C₈alkyl, C₁-C₈alkylthio C₁-C₈alkoxy and/or -N(R₈)₂; or R₃ is -CO-OR₉ or -CO-N(R₉)₂;

Q is a single bond, CR₆R₇, -O- or -S-;

R₄ and R₅ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy;

R₆ and R₇ are each independently of the other hydrogen or C₁-C₄alkyl;

R₈ is C₁-C₁₈ alkyl, C₂-C₁₈ alkyl which is interrupted by one or several non-successive O-atoms; or -N(R₈)₂ forms a 5- or 6-membered O-, S- or N-containing heterocyclic ring;

R₉ is hydrogen, C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O atoms, C₃-C₁₂-cycloalkyl, C₂-C₁₈-alkenyl, phenyl-C₁-C₄-alkyl, phenyl, naphthyl, pyridyl, the radicals phenyl, naphthyl or pyridyl being unsubstituted or substituted by one to five C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio and/or halogen; or -N(R₉)₂ forms a 5- or 6-membered O-, S- or N-containing heterocyclic ring;

R₁₀ is C₁-C₁₈alkyl, C₂-C₁₈alkyl which is interrupted by one or several non-successive O-atoms, C₃-C₁₂-cycloalkyl, phenyl-C₁-C₄-alkyl, C₂-C₁₈-alkenyl, phenyl, naphthyl or biphenyl, the radicals phenyl-C₁-C₄-alkyl, phenyl, naphthyl or biphenyl being unsubstituted or substituted by one to five C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio and/or halogen;

R₁₁ is hydrogen or C₁-C₄-alkyl;

R₁₂ is hydrogen or C₁-C₄-alkyl;

by

(1) reacting diphosphanes of the formula (R₁)₂-P-P(R₁)₂ or polyphosphanes of the formula [R₁P]_n, wherein **n** is ≥ 3 and R₁ is any group as defined for R₁ above

with an alkali metal in a solvent in the presence of a proton source,

(2) followed by the reaction with acid halides of formula III

$$\text{Hal} - \overset{\text{O}}{\parallel} - \text{R}_2 \quad (\text{III}), \text{ and/or by reaction with electrophilic compounds } \text{R}_3\text{-Hal, wherein R}_2, \text{R}_3 \text{ and Hal}$$
 have the meaning cited above.